arXiv:1112.0748v1 [math.CO] 4 Dec 2011

A Note on $\{k, n-k\}$-Factors of Regular Graphs

Hongliang Lu[1] and David G. L. Wang[2]

[1]Department of Mathematics
[1]Xi'an Jiaotong University, Xi'an 710049, P. R. China
[1]`luhongliang215@sina.com`

[2]Beijing International Center for Mathematical Research
[2]Peking University, Beijing 100871, P. R. China
[2]`wgl@math.pku.edu.cn`

Abstract

Let r be an odd integer, and k an even integer. In this note, we present r-regular graphs which have no $\{k, r-k\}$-factors for all $1 \leq k \leq \frac{r}{2}-1$. This gives a negative answer to a problem posed by Akbari and Kano recently.

Keywords: regular graph, $\{k, n-k\}$-factor

2010 AMS Classification: 05C75

1 Introduction

Let G be a simple graph, and H a set of integers. A spanning subgraph F of G is called an *H-factor* if the degree of each vertex belongs to H.

Let r, k be positive integers such that $1 \leq k \leq \frac{r}{2}$. Recently, Akbari and Kano [1] considered whether a general r-regular graph contains a $\{k, r-k\}$-factor. It is known [4] that the problem of determining whether G has an $\{a, b\}$-factor is NP-complete for any positive integers a and b such that $b - a \geq 3$.

The case that both r and k are even was solved by Peterson [6] early in 1891. In fact Peterson proved that any r-regular graph has a $\{k\}$-factor. The proof can be found in Akiyama and Kano's book [1]. For the case that r is odd and k is even, Akbari and Kano [2] showed the existence of a $\{k, r-k\}$-factor of any r-regular graphs by induction on edge-connectivity. In fact, they considered general graphs (allowing multiple edges and loops) and obtained some refined results. They posed the problem of solving the remaining cases.

In this note, we deal with the case that r is even and k is odd.

2 The main result

Let r be an even integer, and k an odd integer, such that $1 \leq k \leq \frac{r}{2}$. In this section, we consider the problem that whether every r-regular graph contains a $\{k, r-k\}$-factor.

First, the case $k = \frac{r}{2}$ can be completely solved by the following result of Gallai [3].

Theorem 2.1 (Gallai) *Every m-edge-connected r-regular graph on an even number of vertices has a k-factor if r is even, k is odd, and*

$$\frac{r}{m} \le k \le r\left(1 - \frac{1}{m}\right).$$

Theorem 2.2 *Let r be an even integer such that $\frac{r}{2}$ is odd. Let G be a connected r-regular graph. Then G has an $\frac{r}{2}$-factor if and only if the number of vertices of G is even.*

Proof. The necessity is clear since $\frac{r}{2}$ is odd. Suppose that G is an r-regular graph with an even number of vertices. Note that any connected r-regular graph is 2-edge-connected if r is even. Taking $m = 2$ in Gallai's Theorem 2.1, we see that G has an $\frac{r}{2}$-factor. $\square$

For the other cases, we will present r-regular graphs without $\{k, r-k\}$-factors. Here is our main result.

Theorem 2.3 *Let r, k be positive integers such that r is even, k is odd, and*

$$1 \le k \le \frac{r}{2} - 1.$$

Then there exists an r-regular graph without $\{k, r-k\}$-factors.

Proof. Note that the case $\frac{r}{2}$ is even and $k = \frac{r}{2} - 1$ is solved by the first author [5]. We divide the remaining cases into two parts depending on the parity of $\frac{r}{2}$. For either of them we shall present an r-regular graph without $\{k, r-k\}$-factors.

Let H be the graph obtained by removing an edge from the complete graph K_{r+1}. We will use H as building blocks in the constructions. Let $H_1, H_2, \ldots$ be pairwise disjoint copies of H. Let u_i and v_i be the unsaturated vertices of H_i.

Suppose that $\frac{r}{2}$ is odd. Then $k \le \frac{r}{2} - 2$. Let G_1 be the graph obtained by linking a vertex v to all unsaturated vertices of H_1, H_2, …, $H_{\frac{r}{2}}$. Then G_1 is an r-regular graph with $r(r+1)/2 + 1$ vertices. Suppose that G_1 contains a $\{k, r-k\}$-factor F_1. By parity, it is easy to deduce that v links to exactly one vertex of each H_i in F_1. It follows that

$$\deg_{F_1}(v) = \frac{r}{2} \notin \{k, r-k\},$$

a contradiction. So G_1 is a desired graph.

Now we consider the case that $\frac{r}{2}$ is even. By the work in [5], we can suppose that $k \le \frac{r}{2} - 3$. In this case, let G_2 be the graph consisting of two vertices u, v, and the copies H_1, H_2, …, H_r such that u is linked to all unsaturated vertices of the copies H_1, H_2, …, $H_{\frac{r}{2}-1}$, and to the vertices u_{r-1} and u_r; while v is linked to all unsaturated vertices of $H_{\frac{r}{2}}$, $H_{\frac{r}{2}+1}$, …, H_{r-2}, and to the vertices v_{r-1} and v_r. Then G_2 is an r-regular graph with $r(r+1)+2$ vertices. Suppose that G_2 contains a $\{k, r-k\}$-factor F_2. By parity, the vertex u links to exactly one vertex of each H_i $(1 \le i \le \frac{r}{2} - 1)$. It follows that

$$\deg_{F_2}(u) \in \left\{\frac{r}{2} - 1,\ \frac{r}{2},\ \frac{r}{2} + 1\right\},$$

which has empty intersection with the set $\{k, r-k\}$. Hence G_2 has no $\{k, r-k\}$-factors. This completes the proof. $\square$

References

[1] S. Akbari and M. Kano, $\{k, r-k\}$-factors of r-regular graphs, to appear.

[2] J. Akiyama and M. Kano, Factors and Factorizations of Graphs, Lecture Notes in Math., vol. 2031, Springer, 2011.

[3] T. Gallai, On factorisation of graphs, Acta Math. Acad. Sci. Hung. 1 (1950), 133–153.

[4] L. Lovász, The factorization of graphs. II, Acta Math. Hungar. 23 (1972), 223–246.

[5] H. Lu, Note on parity factors of regular graphs, arXiv/math.CO: 1110.3198, 2011.

[6] J. Petersen, Die Theorie der regulären graphs, Acta Math. 15 (1891), 193–220.